



## GJB1 gene

gap junction protein beta 1

### Normal Function

The *GJB1* gene provides instructions for making a protein called connexin-32 (also known as gap junction beta 1). This protein is a member of the gap junction connexin family, which plays a role in cell communication by forming channels, or gap junctions, between cells. Gap junctions speed the transport of nutrients, charged particles (ions), and small molecules that carry communication signals between cells.

The connexin-32 protein is made in several tissues, including those of the liver, pancreas, kidney, and nervous system. In the nervous system, this protein is located in the cell membrane of specialized cells called Schwann cells and oligodendrocytes. Schwann cells are found in the peripheral nervous system, which consists of nerves connecting the brain and spinal cord (central nervous system) to muscles and sensory cells that detect sensations such as touch, pain, heat, and sound. Oligodendrocytes are located in the central nervous system.

The Schwann cells and oligodendrocytes surround nerves and are involved in the production and long-term maintenance of a fatty substance called myelin. Myelin forms a protective coating around certain nerve cells and ensures the smooth and rapid transmission of nerve impulses.

The connexin-32 protein forms channels through the myelin sheath, allowing efficient transport and communication between the outer myelin layers and the interior of the Schwann cell or oligodendrocyte.

### Health Conditions Related to Genetic Changes

#### Charcot-Marie-Tooth disease

Researchers have identified approximately 300 *GJB1* gene mutations in people with type X Charcot-Marie-Tooth disease, a disorder characterized by muscle weakness and sensory problems, especially in the hands and feet. A few of these mutations also cause hearing loss in individuals with this type of Charcot-Marie-Tooth disease. Most *GJB1* gene mutations change single protein building blocks (amino acids) in the connexin-32 protein. Some *GJB1* gene mutations result in a protein of abnormal size.

It is unclear how *GJB1* gene mutations lead to the characteristic features of Charcot-Marie-Tooth disease, including a loss of myelin (demyelination) and the slowed transmission of nerve impulses in the peripheral nervous system. The altered protein may be degraded quickly or trapped inside the cell, preventing it from reaching the


cell membrane to form gap junctions. In some cases, an altered protein reaches the cell membrane but does not form properly functioning gap junctions. The loss of functional gap junctions probably impairs the normal activities of Schwann cells, such as myelin production. Malfunctioning gap junctions could also disrupt communication between Schwann cells and the underlying nerve cell, disturbing the transmission of nerve impulses.

In addition to the peripheral nervous system problems associated with this disorder, evidence of demyelination in the central nervous system has been reported in some people with Charcot-Marie-Tooth disease caused by *GJB1* gene mutations. These central nervous system abnormalities did not generally cause any symptoms, but were identified by electrical testing of nerve impulses or imaging studies. Research suggests that another connexin protein whose function overlaps with that of connexin-32 helps compensate for the mutated connexin-32 protein in the oligodendrocytes of the central nervous system.

### Chromosomal Location

Cytogenetic Location: Xq13.1, which is the long (q) arm of the X chromosome at position 13.1

Molecular Location: base pairs 71,215,212 to 71,225,215 on the X chromosome (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

### Other Names for This Gene

- CMTX
- CMTX1
- connexin 32
- CX32
- CXB1\_HUMAN

- gap junction protein, beta 1, 32kDa
- gap junction protein, beta 1, 32kDa (connexin 32, Charcot-Marie-Tooth neuropathy, X-linked)

## **Additional Information & Resources**

### Educational Resources

- Biochemistry (fifth edition, 2002): Gap Junctions Allow Ions and Small Molecules to Flow between Communicating Cells  
<https://www.ncbi.nlm.nih.gov/books/NBK22492/>

### GeneReviews

- Charcot-Marie-Tooth Neuropathy X Type 1  
<https://www.ncbi.nlm.nih.gov/books/NBK1374>

### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28GJB1%5BTIAB%5D%29+OR+%28connexin+32%5BTIAB%5D%29+OR+%28cx+32%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>

### OMIM

- GAP JUNCTION PROTEIN, BETA-1  
<http://omim.org/entry/304040>

### Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology  
[http://atlasgeneticsoncology.org/Genes/GC\\_GJB1.html](http://atlasgeneticsoncology.org/Genes/GC_GJB1.html)
- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=GJB1%5Bgene%5D>
- Connexin-Deafness Homepage  
<http://davinci.crg.es/deafness/>
- HGNC Gene Family: Gap junction proteins  
<http://www.genenames.org/cgi-bin/genefamilies/set/314>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=4283](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=4283)

- Inherited Peripheral Neuropathies Mutation Database  
<http://www.molgen.ua.ac.be/CMTMutations/Mutations/Mutations.cfm?Context=3>
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/2705>
- UniProt  
<http://www.uniprot.org/uniprot/P08034>

## Sources for This Summary

- Abrams CK, Oh S, Ri Y, Bargiello TA. Mutations in connexin 32: the molecular and biophysical bases for the X-linked form of Charcot-Marie-Tooth disease. *Brain Res Brain Res Rev.* 2000 Apr; 32(1):203-14. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/10751671>
- Baker SK, Reith CC, Ainsworth PJ. Novel 95G>A (R32K) somatic mosaic connexin 32 mutation. *Muscle Nerve.* 2008 Nov;38(5):1510-4. doi: 10.1002/mus.21145.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/18949782>
- Kleopa KA, Scherer SS. Molecular genetics of X-linked Charcot-Marie-Tooth disease. *Neuromolecular Med.* 2006;8(1-2):107-22. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/16775370>
- Kochanski A, Kabzinska D. Molecular genetic analysis of the GJB1 gene: a study of six mutations. *J Appl Genet.* 2004;45(1):95-100.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/14960772>
- Mandich P, Grandis M, Geroldi A, Acquaviva M, Varese A, Gulli R, Ciotti P, Bellone E. Gap junction beta 1 (GJB1) gene mutations in Italian patients with X-linked Charcot-Marie-Tooth disease. *J Hum Genet.* 2008;53(6):529-33. doi: 10.1007/s10038-008-0280-4. Epub 2008 Apr 1.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/18379723>
- Mazzeo A, Di Leo R, Toscano A, Muglia M, Patitucci A, Messina C, Vita G. Charcot-Marie-Tooth type X: unusual phenotype of a novel CX32 mutation. *Eur J Neurol.* 2008 Oct;15(10):1140-2. doi: 10.1111/j.1468-1331.2008.02263.x. Epub 2008 Aug 20.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/18717720>
- Ressot C, Bruzzone R. Connexin channels in Schwann cells and the development of the X-linked form of Charcot-Marie-Tooth disease. *Brain Res Brain Res Rev.* 2000 Apr;32(1):192-202. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/10751670>
- Sargiannidou I, Vavlitou N, Aristodemou S, Hadjisavvas A, Kyriacou K, Scherer SS, Kleopa KA. Connexin32 mutations cause loss of function in Schwann cells and oligodendrocytes leading to PNS and CNS myelination defects. *J Neurosci.* 2009 Apr 15;29(15):4736-49. doi: 10.1523/JNEUROSCI.0325-09.2009.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/19369543>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2721059/>
- Wang HL, Chang WT, Yeh TH, Wu T, Chen MS, Wu CY. Functional analysis of connexin-32 mutants associated with X-linked dominant Charcot-Marie-Tooth disease. *Neurobiol Dis.* 2004 Mar; 15(2):361-70.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/15006706>

- Young P, Suter U. The causes of Charcot-Marie-Tooth disease. Cell Mol Life Sci. 2003 Dec;60(12):2547-60. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/14685682>
  - Zambelis T, Panas M, Kokotis P, Karadima G, Kararizou E, Karandreas N. Central motor and sensory pathway involvement in an X-linked Charcot-Marie-Tooth family. Acta Neurol Belg. 2008 Jun;108(2):44-7.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/18795595>
- 

Reprinted from Genetics Home Reference:  
<https://ghr.nlm.nih.gov/gene/GJB1>

Reviewed: August 2009

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services